

species with limited distributions, and the Cameroun-Nigeria border region and western Cameroun contain numerous endemic species. It is therefore likely that *L. priscilla* will be found also in Cameroun. It must be very local and scarce.

The closest relative of the new species, *L. ferrymani*, has quite different ecological preferences. It is one of the few West African Lipteninae found in the Guinea savannah zone; most of the material in The Natural History Museum, London is from as far north as Kaduna. *L. septistrigata*, however, is much more of a forest species.

Acknowledgements

I am grateful to Mr R.G.T. St. Leger for allowing me to describe the new species in this paper, which is no. 12 resulting from preparatory work for the book *Butterflies of West Africa – origins, natural history, diversity, and conservation*. The Carlsberg Foundation in Denmark supports my own field work, for which I am most grateful. I also wish to thank The Natural History Museum, London, and especially Messrs P. Ackery and C.R. Smith for their help and co-operation. The library of the Royal Entomological Society, London, and the assistance of the librarian, Ms Jacqueline Ruffle is also invaluable.

References

- Dowsett, R.J., Hecq, J. & Knoop, D.P., 1989. Ecological notes on two collections of butterflies (Lepidoptera) from eastern Nigeria. In A preliminary natural history survey of Mambilla Plateau and some lowland forests of eastern Nigeria. *Tauraco research Report*, no. 1.
- Libert, M., 1991. Insularité continentale: le cas des Lépidoptères Rhopalocères de la Dorsale camérounaise. *Bulletin de la Société entomologique de France*, **96**: 375-398.
- St. Leger, R.G.T., 1965. The larger butterflies of the Obudu Plateau. *Nigerian Field* **30**: 69-81.
- Stempffer, H., Bennett, N.H. & May, S.J., 1974. A revision of some groups of *Liptena* Westwood (Lepidoptera: Lycaenidae). *Bulletin of the British Museum, Natural History (Entomology)*, **30**: 109-181 (+plates).

Euchmichtis lichenea lichenea (Hb.) Feathered *Ranunculus* (Lep.: Noctuidae) in north-east Hampshire

At about 10pm on 8th October 1994 a male specimen of *Euchmichtis lichenea lichenea* flew to light at the cottage here in Selborne. This occurrence in north-east Hampshire would seem to represent an extension of *lichenea*'s previously published range in the south and north-west of the county.

References: Bernard Skinner & David Wilson, 1984. *Colour Identification Guide to Moths of the British Isles* Viking, Harmondsworth; Heath, J. & Emmet, A.M. (eds.), 1983. *The Moths and Butterflies of Great Britain and Ireland* **10** Harley Books, Colchester; Goater, B., 1974. *The Butterflies & Moths of Hampshire and the Isle of Wight*, E.W. Classey, Faringdon; Goater, B., 1992. *The Butterflies and Moths of Hampshire and the Isle of Wight: additions and corrections*, Joint Nature Conservation Committee, Peterborough.

– ALASDAIR ASTON, Wake's Cottage, Selborne, Hampshire GU34 3JH.